

JONES POND

2018 SAMPLING HIGHLIGHTS

Station – 3 Deep

New Durham, NH



Blue = Oligotrophic

Yellow = Mesotrophic

Red = Eutrophic

Gray = No Data




Figure 1. Jones Pond Water Quality (2018)

Table 1. 2018 Jones Pond Seasonal Averages and NH DES Aquatic Life Nutrient Criteria¹

Parameter	Oligotrophic	Mesotrophic	Eutrophic	Jones Pond Average (range)	Jones Pond Classification
Water Clarity (meters)	4.0 – 7.0	2.5 - 4.0	< 2.5	3.1 meters (2.2 – 3.8)	Mesotrophic
Chlorophyll <i>a</i> ¹ (ug/l)	< 3.3	> 3.3 – 5.0	> 5.0 – 11.0	8.0 ug/l (1.9 – 17.2)	Eutrophic
Total Phosphorus ¹ (ug/l)	< 8.0	> 8.0 – 12.0	> 12.0 – 28.0	26.2 ug/l (15.8 – 35.8)	Eutrophic
Dissolved Oxygen (mg/L)	5.0 – 7.0	2.0 – 5.0	<2.0	* 0.3 mg/l (0.1 – 0.6)	* Eutrophic

* Jones Pond developed a mid-lake layer of rapidly decreasing temperature and the August 9 oxygen profile was used to classify Jones Pond.

Table 2. 2018 Jones Pond Seasonal Average Accessory Water Quality Measurements

Parameter	Assessment Criteria					Jones Pond Average (range)	Jones Pond Classification
Color (color units)	< 10 uncolored	10 – 20 slightly colored	20 – 40 lightly tea colored	40 – 80 tea colored	> 80 highly colored	40.4 color units (range: 25.9 – 84.1)	Tea colored
Alkalinity (mg/L)	< 0.0 acidified	0.1 – 2.0 extremely vulnerable	2.1 – 10 moderately vulnerable	10.1 – 25.0 low vulnerability	> 25.0 not vulnerable	7.5 mg/L (range: 5.8 – 8.2)	Moderately vulnerable
pH (std units)	< 5.5 suboptimal for successful growth and reproduction		6.5 – 9.0 optimal range for fish growth and reproduction			6.7 standard units (range: 6.4 – 6.9)	Optimal range for fish growth and reproduction
Specific Conductivity (uS/cm)	< 50 uS/cm Characteristic of minimally impacted NH lakes		50-100 uS/cm Lakes with some human influence	> 100 uS/cm Characteristic of lakes experiencing human disturbances		56.4 uS/cm (range: 45.0 – 61.1)	Characteristic of lakes with some human influence






Figure 2 and 3. Seasonal Secchi disk transparency, chlorophyll *a* concentrations and dissolved color concentrations. Figures 2 and 3 illustrate the interplay among Secchi Disk transparency, chlorophyll *a* and dissolved color concentrations. Shallower water transparency measurements oftentimes correspond to increases in chlorophyll *a* and/or color concentrations.

Table 3. Merrymeeting River watershed inter-lake comparison (2018 Data)

Lake	Average (range) Secchi Disk Transparency (meters)	Average (range) Chlorophyll <i>a</i> (ppb)	Average (range) Total Phosphorus (ppb)	Average (range) Dissolved Color (CPU)	Average (range) Dissolved Oxygen (mg/l)
Merrymeeting Lake	10.8 meters (range: 8.6 – 12.1)	0.6 ug/l (range: 0.3 – 0.9)	3.4 ug/l (range: 2.0 – 5.2)	4.6 CPU (range: 2.7 – 6.3)	11.1 mg/l (range: 8.1 – 13.1)
Marsh Pond	3.4 meters (range: 2.5 – 5.2)	5.7 ug/l (range: 1.6 – 18.8)	28.2 ug/l (range: 14.1 – 46.8)	34.1 CPU (range: 19.3 – 72.4)	2.7 mg/l (range: 0.2 – 9.2)
Jones Pond	3.1 meters (range: 2.2 – 3.8)	8.0 ug/l (range: 1.9 – 17.2)	26.2 ug/l (range: 15.8 – 35.8)	40.4 CPU (range: 25.9 – 84.1)	0.3 mg/l (range: 0.1 – 0.6)
Downing Pond	3.0 meters (range: 2.7 – 3.3)	5.1 ug/l (range: 1.7 – 8.9)	24.6 ug/l (range: 16.0 – 34.3)	40.1 CPU (range: 29.5 – 74.2)	-----

- Water quality data are reported for a deep reference sampling location in each lake/pond.
- Dissolved oxygen measurements were collected in the summer (late-July and mid-August) in the bottom water layer (hypolimnion or metalimnion).
- Downing Pond Secchi Disk transparency measurements intermittently reached the lake bottom before disappearing from view and likely underestimate the water transparency.
- ----- Indicates the site is too shallow to form a stable deep water layer (hypolimnion or metalimnion) during the summer months.

Figure 4. Jones Pond dissolved oxygen profile collected between June 7 and November 6, 2018. The vertical red line indicates the oxygen concentration commonly considered the threshold for successful growth and reproduction of warm water fish such as bass and perch.



Recommendations


Implement Best Management Practices within the Jones Pond watershed to minimize the adverse impacts of polluted runoff and erosion on Jones Pond. Refer to “Landscaping at the Water’s Edge: An Ecological Approach” and “New Hampshire Homeowner’s Guide to Stormwater Management: Do-It-Yourself Stormwater Solutions for Your Home” for more information on how to reduce nutrient loading caused by overland run-off.

- https://extension.unh.edu/resources/files/Resource004159_Rep5940.pdf
- <https://des.nh.gov/organization/commissioner/pip/publications/wd/documents/wd-11-11.pdf>

A Merrymeeting River watershed management plan is in development. Once completed, the Watershed Management Plan will identify threats to Jones Pond and will provide potential mitigation strategies that will help stabilize and improve water quality.

Figure 5. Jones Pond

New Durham, NH
2018 Deep water sampling site



0 0.1 0.2 0.3 0.4 Miles

Aerial Orthophoto Source: NH GRANIT
GPS Coordinates collected by the UNH Center for Freshwater Biology



Extension

